## **Triadobatrachus**

**Triadobatrachus** is an extinct genus of <u>salientian</u> <u>frog-like</u> amphibians, including only one known species, *Triadobatrachus massinoti*. It is the oldest member of the frog lineage known, and an excellent example of a <u>transitional fossil</u>. It lived during the <u>Early Triassic</u> about 250 million years ago, in what is now <u>Madagascar</u>.



Life restoration by Pavel Riha

**Triadobatrachus** was 10 cm (3.9 in) long, and retained still primitive characteristics, such as possessing at least vertebrae. modern frogs have only four to nine. At least 10 of these vertebrae formed a short tail, which the animal may have retained as an adult.<sup>[1]</sup> It probably swam by kicking its hind legs, although it could not jump,

as most modern frogs can. Its  $\underline{\text{skull}}$  resembled that of modern frogs, consisting of a latticework of thin bones separated by large openings.<sup>[2]</sup>

This creature, or a relative, evolved eventually into modern frogs, the earliest example of which is *Prosalirus*, millions of years later in the Early Jurassic.<sup>[3]</sup>


It was first discovered in the 1930s, when <u>Adrien Massinot</u>, near the village of <u>Betsiaka</u> in northern Madagascar, found an almost complete skeleton in the <u>Induan Middle Sakamena Formation</u> of the <u>Sakamena Group</u>. The animal must have fossilized soon after its death, because all bones lay in their natural anatomical position.

**Triadobatrachus** Temporal range: Early Triassic, 250 Ma Pre€ € OS D C P T J K PgN Slabs of the fossil Scientific classification 🥖 Kingdom: Animalia Phylum: Chordata Class: **Amphibia** Genus: †Triadobatrachus Kuhn, 1962 Species: †T. massinoti **Binomial** name †Triadobatrachus massinoti (Piveteau, 1936)

Only the anterior part of the skull and the ends of the limbs were missing. This fossil was initially described under the name *Protobatrachus massinoti* by the French paleontologist Jean Piveteau in 1936.<sup>[4][5]</sup> Much more detailed description were published more recently.<sup>[1][6]</sup>

Although it was found in marine deposits, the general structure of *Triadobatrachus* shows that it probably lived for part of the time on land and breathed air. Its proximity to the mainland is further borne out by the remains of terrestrial plants found with it, and because most extant amphibians do not tolerate saltwater,<sup>[7]</sup> and that this saltwater intolerance was probably present in the earliest <u>lissamphibians</u>.<sup>[8]</sup>

## **Gallery**









CT-scan

Diagram

Cast

Life restoration by N. Tamura

## References

- Ascarrunz, Eduardo; Rage, Jean-Claude; Legreneur, Pierre; Laurin, Michel (2016).
  "Triadobatrachus massinoti, the earliest known lissamphibian (Vertebrata: Tetrapoda) reexamined by μCT-Scan, and the evolution of trunk length in batrachians" (http://repository.naturalis.nl/cgi/b/bib/bib-idx?c=naturalis;type=simple;view=reslist;fmt=long;rgn1=id;q1=oai%3AARN O%3A610011;lang=en;sort=year;cc=naturalis;page=reslist;size=1;start=1). Contributions to Zoology. 58 (2): 201–234.
- 2. Palmer, D., ed. (1999). *The Marshall Illustrated Encyclopedia of Dinosaurs and Prehistoric Animals*. London: Marshall Editions. p. 56. ISBN 1-84028-152-9.
- 3. Rocek, Z. (2000). "14. Mesozoic Amphibians". In Heatwole, H.; Carroll, R. L. (eds.). <u>Amphibian Biology. Paleontology: The Evolutionary History of Amphibians</u> (http://rocek.gli.cas.cz/Reprints/AmphBiol3.pdf) (PDF). **4.** Surrey Beatty & Sons. pp. 1295–1331. ISBN 0-949324-87-6.
- 4. Piveteau, J. (1936). "Une forme ancestrale des amphibiens anoures dans le Trias inférieur de Madagascar". *Comptes Rendus Hebdomadaires des Séances de l'Académie des Sciences*. **202**: 1607–1608.
- 5. Piveteau, J. (1936). "Origine et évolution morphologique des amphibiens anoures". *Comptes Rendus Hebdomadaires des Séances de l'Académie des Sciences*. **203**: 1084–1086.
- 6. Rage, J-C; Roček, Z. (1989). "Redescription of *Triadobatrachus massinoti* (Piveteau, 1936) an anuran amphibian from the Early Triassic". *Palaeontographica Abteilung A.* **206**: 1–16.
- 7. Hopkins, G. R.; Brodie Jr, E. D. (2015). "Occurrence of Amphibians in Saline Habitats: A Review and Evolutionary Perspective" (http://www.hljournals.org/doi/abs/10.1655/HERPMON OGRAPHS-D-14-00006). Herpetological Monographs. 29 (1): 1–27. doi:10.1655/herpmonographs-d-14-00006 (https://doi.org/10.1655%2Fherpmonographs-d-14-00006).
- 8. Laurin, Michel; Soler-Gijón, Rodrigo (2010). "Osmotic tolerance and habitat of early stegocephalians: indirect evidence from parsimony, taphonomy, paleobiogeography, physiology and morphology" (http://sp.lyellcollection.org/content/339/1/151.short). *Geological Society of London, Special Publications*. **339**: 151–179. doi:10.1144/sp339.13 (https://doi.org/10.1144%2Fsp339.13).
- Benes, Josef. Prehistoric Animals and Plants. Pg. 114. Prague: Artia, 1979.

Retrieved from "https://en.wikipedia.org/w/index.php?title=Triadobatrachus&oldid=946402080"

This page was last edited on 19 March 2020, at 22:56 (UTC).

Text is available under the <u>Creative Commons Attribution-ShareAlike License</u>; additional terms may apply. By using this site, you agree to the Terms of Use and Privacy Policy. Wikipedia® is a registered trademark of the Wikimedia



Foundation, Inc., a non-profit organization.